

MULTIFRACTALITY OF WHOLE-PLANE SLE

Bertrand Duplantier*, **Michel Zinsmeister****

Nguyen T.P. Chi**, **Nguyen T.T. Nga****

*Institut de Physique Théorique, Saclay, France

**MAPMO, Université d'Orléans, France

COLLOQUE GDR MULTIFRACTALES 2013

Île de Porquerolles

22-27 septembre 2013

Let $f(z) = \sum_{n \geq 0} a_n z^n$ be a holomorphic function in the unit disc \mathbb{D} . Further assume that f is injective. Then $a_1 \neq 0$ and *Bieberbach* proved in 1916 that $|a_2| \leq 2|a_1|$. In the same paper, he famously conjectured that $\forall n \geq 2, |a_n| \leq n|a_1|$, guided by the intuition that the *Koebe function*

$$\mathcal{K}(z) := - \sum_{n \geq 1} n(-z)^n = \frac{z}{(1+z)^2},$$

which is a holomorphic bijection between \mathbb{D} and $\mathbb{C} \setminus [1/4, +\infty)$, should be *extremal*. This conjecture was finally proven in 1984 by *de Branges*. The earliest important contribution to the proof of Bieberbach's conjecture is that by *Loewner* in 1923 that $|a_3| \leq 3|a_1|$. *Oded Schramm* revived Loewner's method in 1999, introducing *randomness* into it, as driven by *standard Brownian motion*.

Whole-Plane SLE & LLE

$$\frac{\partial f_t}{\partial t} = z \frac{\partial f_t}{\partial z} \frac{\lambda(t) + z}{\lambda(t) - z}, \quad z \in \mathbb{D},$$

$$\lambda(t) = e^{i\sqrt{\kappa}B_t} [e^{i\xi L_t}].$$

The characteristic function of a Lévy process L_t has the form


$$\mathbb{E}(e^{i\xi L_t}) = e^{-t\eta(\xi)},$$

where η the Lévy symbol. The function

$$\eta(\xi) = \kappa|\xi|^\alpha/2, \quad \alpha \in (0, 2]$$

is the Lévy symbol of the α -stable process. The normalization here is chosen so that it is SLE_κ for $\alpha = 2$.

Whole-Plane SLE ($\mathbb{D} \rightarrow \mathbb{C} \setminus \gamma[0, \infty)$)



Loewner map $z \mapsto f_t(z)$ from the unit disk \mathbb{D} to the slit domain $\Omega_t = \mathbb{C} \setminus \gamma([t, \infty))$. One has $f_t(0) = 0, \forall t \geq 0$. At $t = 0$, $\lambda(0) = 1$, so that the image of $z = 1$ is at the tip $\gamma(0) = f_0(1)$ of the curve.

Series expansions

Let f_t be the whole-plane evolution generated by the Lévy process (L_t) with Lévy symbol η . We write

$$e^{-t} f_t(z) = z + \sum_{n=2}^{\infty} a_n(t) z^n; \quad e^{-t/2} h_t(z) = z + \sum_{n \geq 1} b_{2n+1}(t) z^{2n+1}.$$

Then the *conjugate* whole-plane LLE $e^{-iL_t} f_t(e^{iL_t} z)$ has the same law as $f_0(z)$, i.e., $e^{i(n-1)L_t} a_n(t) \stackrel{(\text{law})}{=} a_n(0)$. Similarly, the *conjugate of the oddified* whole-plane LLE $h_t(z) := z \sqrt{f_t(z^2)/z^2}$, $e^{-(i/2)L_t} h_t(e^{(i/2)L_t} z)$, has the same law as $h_0(z)$, i.e., $e^{inL_t} b_n(t) \stackrel{(\text{law})}{=} b_n(0)$.

Loewner's method

Recall that

$$f_t(z) = e^t \left(z + \sum_{n \geq 2} a_n(t) z^n \right).$$

By expanding both sides of Loewner's equation as power series, and identifying coefficients, leads one to the set of *recursion equations for*
 $n \geq 2$

$$\dot{a}_n(t) - (n-1)a_n(t) = 2 \sum_{k=1}^{n-1} k a_k(t) \bar{\lambda}^{n-k}(t),$$

with $a_1 = 1$; *the dot means a t -derivative*, and $\bar{\lambda}(t) = 1/\lambda(t)$, with
 $\lambda(t) = e^{i\sqrt{\kappa}B_t} [e^{i\xi L_t}]$.

Expected coefficients

Theorem 1. *Setting $a_n := a_n(0)$ and $b_{2n+1} := b_{2n+1}(0)$, we have*

$$\mathbb{E}(a_n) = \prod_{k=0}^{n-2} \frac{\eta_k - k - 2}{\eta_{k+1} + k + 1}, \quad n \geq 2,$$

$$\mathbb{E}(b_{2n+1}) = \prod_{k=0}^{n-1} \frac{\eta_k - k - 1}{\eta_{k+1} + k + 1}, \quad n \geq 1.$$

Corollary 1. *If $\eta_1 = 3$, $\mathbb{E}(f'_0(z)) = 1 - z$ (SLE_6);*


if $\eta_1 = 1$ and $\eta_2 = 4$, $\mathbb{E}(f'_0(z)) = (1 - z)^2$ (SLE_2);

if $\eta_1 = 2$, $\mathbb{E}(h'_0(z)) = 1 - z^2$ (SLE_4).

[See also [Kemppainen '10](#) for expectations of SLE coefficient moments.]

The Surprise: Expected Square Coefficients $\mathbb{E}(|a_n|^2)$

Example: For SLE_6



Expected Square Coefficients

Example: For SLE_6

$$\mathbb{E}(|a_n|^2) = 1, \kappa = 6, \forall n$$

$$\mathbb{E}(|a_4|^2) = \frac{8\kappa^5 + 104\kappa^4 + 4576\kappa^3 + 18288\kappa^2 + 22896\kappa + 8640}{9(\kappa + 10)(3\kappa + 2)(\kappa + 6)(\kappa + 1)(\kappa + 2)^2}.$$

[Recursion: $n \leq 4$; Computer assisted: $n \leq 8$ (formal), $n \leq 19$ (num.)]

Theorem 2.

- (i) if $\eta_1 = 3$, $\mathbb{E}(|a_n|^2) = 1, \forall n \geq 1$ (SLE_6);
- (ii) if $\eta_1 = 1, \eta_2 = 4$, $\mathbb{E}(|a_n|^2) = n, n \geq 1$ (SLE_2);
- (iii) if $\eta_1 = 2$, $\mathbb{E}(|b_{2n+1}|^2) = 1/(2n+1), n \geq 1$ (SLE_4).

Derivative Moments

Theorem 3. *The whole-plane SLE_{κ} map $f_0(z)$ has derivative moments*

$$\begin{aligned}\mathbb{E}[(f'_0(z))^{p/2}] &= (1-z)^{\alpha}, \\ \mathbb{E}[|f'_0(z)|^p] &= \frac{(1-z)^{\alpha}(1-\bar{z})^{\alpha}}{(1-z\bar{z})^{\beta}},\end{aligned}$$

*for the special set of exponents $p = p(\kappa) := (6 + \kappa)(2 + \kappa)/8\kappa$, with $\alpha = (6 + \kappa)/2\kappa$ and $\beta = (6 + \kappa)^2/8\kappa$. [See also *Loutsenko & Yermolayeva '12*]*

Corollary 4. *$p = 2$ case: for $\kappa = 6$:*

$$\mathbb{E}(f'_0(z)) = 1 - z, \quad \mathbb{E}(|f'_0(z)|^2) = \frac{(1-z)(1-\bar{z})}{(1-z\bar{z})^3};$$

for $\kappa = 2$:

$$\mathbb{E}(f'_0(z)) = (1-z)^2, \quad \mathbb{E}(|f'_0(z)|^2) = \frac{(1-z)^2(1-\bar{z})^2}{(1-z\bar{z})^4}.$$

The BS Equation

Beliaev and Smirnov (2005) obtained by martingale arguments the following equation for the *exterior whole-plane* case

$$(F(z) = F(re^{i\theta}), r \geq 1, \sigma = +1)$$

$$p \left(\frac{r^4 + 4r^2(1 - r \cos \theta) - 1}{(r^2 - 2r \cos \theta + 1)^2} - \sigma \right) F + \frac{r(r^2 - 1)}{r^2 - 2r \cos \theta + 1} F_r - \frac{2r \sin \theta}{r^2 - 2r \cos \theta + 1} F_\theta + \Lambda F = 0.$$

Proposition 1. *For the interior whole-plane Schramm (or Lévy)-Loewner evolution, the moments of the derivative modulus, $F(z) := \mathbb{E}(|f'_0(z)|^p)$, satisfy the same BS equation, but with $\sigma = -1$, and $\Lambda = (\kappa/2)\partial^2/\partial\theta^2$ the generator of the driving Brownian process (or of the Lévy process).*

Holomorphic Coordinates

Switch to z, \bar{z} variables, instead of polar coordinates, and write $F(z)$ above as

$$F(z, \bar{z}) := \mathbb{E}(|f'_0(z)|^p) = \mathbb{E}[(f'_0(z))^{p/2}(\bar{f}'_0(\bar{z}))^{p/2}].$$

Using $\partial := \partial_z, \bar{\partial} := \partial_{\bar{z}}$, the equation then becomes

$$-\frac{\kappa}{2}(z\partial - \bar{z}\bar{\partial})^2 F + \frac{z+1}{z-1}z\partial F + \frac{\bar{z}+1}{\bar{z}-1}\bar{z}\bar{\partial} F - p \left[\frac{1}{(z-1)^2} + \frac{1}{(\bar{z}-1)^2} + (\sigma - 1) \right] F = 0.$$

Exterior/Interior whole-plane: $\sigma = \pm 1$.

The action of the differential operator $\mathcal{P}(D)$ above on a function of the factorized form $F(z, \bar{z}) = \varphi(z)\bar{\varphi}(\bar{z})P(z, \bar{z})$ is, by Leibniz's rule, given by

$$\begin{aligned} \mathcal{P}(D)[\varphi\bar{\varphi}P] = & - \frac{\kappa}{2}\varphi\bar{\varphi}(z\partial - \bar{z}\bar{\partial})^2P - \kappa(z\partial - \bar{z}\bar{\partial})(\varphi\bar{\varphi})(z\partial - \bar{z}\bar{\partial})P \\ & + \kappa(z\partial\varphi)(\bar{z}\bar{\partial}\bar{\varphi})P + \varphi\bar{\varphi}\frac{z+1}{z-1}z\partial P + \varphi\bar{\varphi}\frac{\bar{z}+1}{\bar{z}-1}\bar{z}\bar{\partial}P \\ & + \left[-\frac{\kappa}{2}\bar{\varphi}(z\partial)^2\varphi - \frac{\kappa}{2}\varphi(\bar{z}\bar{\partial})^2\bar{\varphi} + \bar{\varphi}\frac{z+1}{z-1}z\partial\varphi + \varphi\frac{\bar{z}+1}{\bar{z}-1}\bar{z}\bar{\partial}\bar{\varphi} \right] P \\ & - p \left[\frac{1}{(z-1)^2} + \frac{1}{(\bar{z}-1)^2} + \sigma - 1 \right] \varphi\bar{\varphi}P. \end{aligned}$$

- For the particular choice of a rotationally invariant $P(z, \bar{z}) := P(z\bar{z})$, the first line above vanishes.
- Study the algebra generated by the action of $\mathcal{P}(D)$ on $\varphi(z) = \varphi_\alpha(z) := (1-z)^\alpha$, and $P(z\bar{z}) := (1-z\bar{z})^{-\beta}$, $\forall \alpha, \beta$.

Integral means spectrum

Definition 1. *The integral means spectrum of a conformal mapping f is the function defined on \mathbb{R} by*

$$\beta(p) := \overline{\lim}_{r \rightarrow 1} \frac{\log(\int_{\partial D} |f'(rz)|^p |dz|)}{\log(\frac{1}{1-r})}.$$

In the *stochastic* setting, one defines the *average* integral means spectrum

Definition 2.

$$\beta(p) := \overline{\lim}_{r \rightarrow 1} \frac{\log(\int_{\partial D} \mathbb{E} |f'(rz)|^p |dz|)}{\log(\frac{1}{1-r})}.$$

Corollary 5. *For a Lévy-Loewner evolution with $\eta_1 = 1, \eta_2 = 4$, or $\eta_1 = 3$ (thus including SLE for $\kappa = 2, 6$), and for an oddified LLE with $\eta_1 = 2$ (thus including SLE for $\kappa = 4$), one has, respectively:*

$$\mathbb{E} \left(\frac{1}{2\pi} \int_0^{2\pi} |f'(re^{i\theta})|^2 d\theta \right) = \frac{1 + 4r^2 + r^4}{(1 - r^2)^4}; \frac{1 + r^2}{(1 - r^2)^3}; \frac{1 + r^4}{(1 - r^4)^2}.$$

This gives the values of the average integral means spectrum $\beta(2) = 4, 3$ for whole-plane LLE with $\eta_1 = 1, \eta_2 = 4$ or $\eta_1 = 3$ (thus whole-plane SLE with $\kappa = 2, 6$) respectively. For the oddified LLE with $\eta_1 = 2$ (thus the oddified whole-plane SLE₄), $\beta_2(2) = 2$.

- They differ from the corresponding values at $p = 2$ of the SLE integral mean spectrum of Beliaev and Smirnov '05.

Define

$$\beta_0(p, \kappa) := -p + \frac{4 + \kappa}{4\kappa} \left(4 + \kappa - \sqrt{(4 + \kappa)^2 - 8\kappa p} \right),$$

$$\hat{\beta}_0(p, \kappa) := p - \frac{(4 + \kappa)^2}{16\kappa}.$$

This is the *average* integral means spectrum $\bar{\beta}_0(p, \kappa)$ of the bulk of SLE_κ , as obtained in *Beliaev & Smirnov '05*:

$$\bar{\beta}_0(p, \kappa) = \beta_0(p, \kappa), \quad 0 \leq p \leq p_0^*(\kappa),$$

$$= \hat{\beta}_0(p, \kappa), \quad p \geq p_0^*(\kappa),$$

$$p_0^*(\kappa) := \frac{3(4 + \kappa)^2}{32\kappa}.$$

Integral means spectra

The whole-plane SLE $_{\kappa}$, $f_{t=0}(z)$, $z \in \mathbb{D}$, and its m -fold transforms, $h_0^{(m)}(z) := z[f_0(z^m)/z^m]^{1/m}$, $m \geq 1$, have average integral means spectra $\beta_m(p, \kappa)$ that exhibit a *phase transition* and are given, for $p \geq 0$, by

$$\beta_1(p, \kappa) = \max \left\{ \beta_0(p, \kappa), 3p - \frac{1}{2} - \frac{1}{2} \sqrt{1 + 2\kappa p} \right\},$$

$$\beta_2(p, \kappa) = \max \left\{ \beta_0(p, \kappa), 2p - \frac{1}{2} - \frac{1}{2} \sqrt{1 + \kappa p} \right\},$$

$$\beta_m(p, \kappa) = \max \left\{ \bar{\beta}_0(p, \kappa), (1 + 2/m)p - \frac{1}{2} - \frac{1}{2} \sqrt{1 + 2\kappa p/m} \right\}.$$

The first spectrum β_1 has its transition point at

$$p^*(\kappa) := \frac{1}{16\kappa} \left((4 + \kappa)^2 - 4 - 2\sqrt{4 + 2(4 + \kappa)^2} \right) < p_0^*(\kappa).$$

Theorem 5. *The average integral means spectrum $\beta(p, \kappa)$ of the unbounded whole-plane SLE_κ has a phase transition at $p^*(\kappa)$ and a special point at $p(\kappa) := (6 + \kappa)(2 + \kappa)/8\kappa$, such that*

$$\beta(p, \kappa) = \beta_0(p, \kappa), \quad 0 \leq p \leq p^*(\kappa);$$

$$\beta(p, \kappa) = 3p - \frac{1}{2} - \frac{1}{2} \sqrt{1 + 2\kappa p} > \beta_0(p, \kappa), \quad p^*(\kappa) < p < p(\kappa);$$

$$\beta(p(\kappa), \kappa) = \frac{(6 + \kappa)^2}{8\kappa};$$

$$\beta(p, \kappa) \leq 3p - \frac{1}{2} - \frac{1}{2} \sqrt{1 + 2\kappa p}, \quad p(\kappa) < p.$$

- For $p > p^*(\kappa)$ the BS solution ceases to be uniformly positive.
- Existence of a *subsolution/supersolution* for the parabolic operator $\mathcal{P}(D)[\psi(z, \bar{z})\ell_\delta(z\bar{z})] \stackrel{\leq}{\geq} 0$ in some annulus of \mathbb{D} whose boundary includes $\partial\mathbb{D}$, corresponding respectively to $p \stackrel{\leq}{\geq} p^*(\kappa)$. Trial functions: $\psi(z, \bar{z}) := (1 - z\bar{z})^{-\beta} |1 - z|^{2\alpha}$, $\ell_\delta(z\bar{z}) := [-\log(1 - z\bar{z})]^\delta$.

Integral means spectrum: *Inner whole-plane SLE*




Figure 1: $\beta(p) = 3p - \frac{1}{2} - \frac{1}{2}\sqrt{1 + 2\kappa p}$

Integral means spectrum: *Outer whole-plane SLE*




Figure 2: $\beta(p) = -p - \frac{1}{2} - \frac{1}{2}\sqrt{1 - 2\kappa p}$, $p^*(\kappa) = (4 + \kappa)^2(8 + \kappa)/128$ (Beliaev, B.D., Zinsmeister '13)

Packing Spectrum

The *packing spectrum* [Makarov] is defined as

$$s(p) := \beta(p) - p + 1.$$

For the unbounded whole-plane SLE_κ , we have for $p \geq p^*(\kappa)$

$$\begin{aligned} s(p, \kappa) &= \beta(p, \kappa) - p + 1 \\ &= 2p + \frac{1}{2} - \frac{1}{2} \sqrt{1 + 2\kappa p}. \end{aligned}$$

Consider its *inverse function*

$$\begin{aligned} p = p(s, \kappa) &:= \frac{s}{2} + \frac{\kappa}{8} u_\kappa^{-1}(s), \\ u_\kappa^{-1}(s) &:= \frac{1}{2\kappa} \left(\kappa - 4 + \sqrt{(4 - \kappa)^2 + 16\kappa s} \right) \end{aligned}$$

(KPZ formula)


Relation to Tip & Derivative Exponents

(Non-standard) tip multifractal exponents obtained by quantum gravity [D. '00], corresponding geometrically to the extremity of an SLE_{κ} path avoiding a packet of s independent Brownian motions.

Differ from the ones associated to the *standard SLE tip multifractal spectrum* [Hastings '02, Beliaev & Smirnov '05, Johansson & Lawler '09].

Identical to the *derivative exponents* obtained for radial SLE_{κ} [Lawler, Schramm & Werner '01].

(Inverse) Radial SLE Map



$$f_0(z) \stackrel{(\text{law})}{=} \lim_{t \rightarrow +\infty} [e^t g_t^{-1}(z) =: \tilde{f}_t(z)].$$

Harmonic measure




Figure 3: $f_0(z) \stackrel{(\text{law})}{=} \lim_{t \rightarrow +\infty} e^t g_t^{-1}(z)$, where $z \mapsto g_t^{-1}(z)$ maps \mathbb{D} to the slit domain $\mathbb{D} \setminus K_t$ (K_t SLE hull). The length $L_t := |g_t(A_t)|$ of the image of the boundary set $A_t := \partial\mathbb{D} \setminus \overline{K_t}$ is the $(2\pi) \times$ the **harmonic measure** of A_t as seen from 0 in $\mathbb{D} \setminus K_t$, with $\mathbb{E}[L_t^s] \asymp e^{-p(s, \kappa)t}$ for $t \rightarrow +\infty$ [LSW '01].

Derivative exponents

Lemma 1. (Lawler, Schramm, Werner '01) Let

$$A_t := \partial\mathbb{D} \setminus \overline{K}_t,$$

which is either an arc on $\partial\mathbb{D}$ or $A_t = \emptyset$. Let $s \geq 0$, and set

$$p = p(s, \kappa) := \frac{s}{2} + \frac{1}{16} \left(\kappa - 4 + \sqrt{(4 - \kappa)^2 + 16\kappa s} \right).$$

Let $\mathcal{H}(\theta, t)$ denote the event $\{w = \exp(i\theta) \in A_t\}$, and set

$$\mathcal{F}(\theta, t) := \mathbb{E} \left[\left| g'_t(\exp(i\theta)) \right|^s 1_{\mathcal{H}(\theta, t)} \right],$$

$$q = q(s, \kappa) := \mathcal{U}_\kappa^{-1}(s) = \frac{\kappa - 4 + \sqrt{(4 - \kappa)^2 + 16\kappa s}}{2\kappa},$$

$$\mathcal{F}(\theta, t) \asymp \exp(-pt) (\sin(\theta/2))^q, \quad \forall t \geq 1, \quad \forall \theta \in (0, 2\pi).$$

Packing spectrum & derivative exponents

The average integral means spectrum involves evaluating, for the whole-plane SLE map $f_0(z)$, the integral

$$\mathbb{I}_p(r) := \int_{\partial D} \mathbb{E} [|f'_0(rz)|^p] |dz|,$$

on a circle of radius $r < 1$ concentric to $\partial\mathbb{D}$, and looking for the smallest $\beta(p)$ such that

$$(1-r)^{\beta(p)} \mathbb{I}_p(r) \stackrel{r \rightarrow 1}{<} +\infty.$$

For $p \geq p^(\kappa)$, the integrand behaves like a distribution and the circle integral concentrates in the vicinity of the pre-image point of infinity by the whole-plane map, $z_0 := f_0^{-1}(\infty) \in \partial\mathbb{D}$. In the large- t approximation to f_0 , that is the neighborhood of $g_t(A_t)$.*

Condensation

The circle integral there is the *restricted* integral in the image w -unit circle

$$I_p(t) := \int_{A_t} e^{pt} |g'_t(w)|^s |dw|; \quad s = s(p) = \beta(p) + 1 - p,$$

From **LSW**'s Lemma above

$$\mathbb{E}[I_p(t)] \asymp \int_0^{2\pi} \sin^q(\theta/2) d\theta < +\infty.$$

By defining the *stochastic radius* $r_t := 1 - L_t \rightarrow 0$, this can be recast as

$$\mathbb{E} \left[(1 - r_t)^{\beta(p)} \int_{\partial\mathbb{D}} |\tilde{f}'_t(r_t z)|^p |dz| \right] \asymp 1, \quad t \rightarrow +\infty,$$

where $f_0(z) \stackrel{(\text{law})}{=} \lim_{t \rightarrow +\infty} [\tilde{f}_t(z) := e^t g_t^{-1}(z)]$. This is (*formally*) reminiscent of the definition of the average integral means spectrum, hinting at why *the derivative exponent* $p = p(s, \kappa)$ *is the inverse function of the unbounded whole-plane packing spectrum* $s(p, \kappa)$. □